

## Guidelines for Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents


Downloaded from <a href="http://aidsinfo.nih.gov/guidelines">http://aidsinfo.nih.gov/guidelines</a> on 12/21/2016

Visit the AIDS*info* website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <a href="http://aidsinfo.nih.gov/e-news">http://aidsinfo.nih.gov/e-news</a>.

## Figure 1 (Last updated May 7, 2013; last reviewed May 7, 2013)

## Immunization Schedule for Human Immunodeficiency Virus (HIV)-Infected Adults



<sup>\*</sup>Covered by the Vaccine Injury Compensation Program

For all persons in this category who meet the age requirements and who lack documentation of vaccination or have no evidence of previous infection;
zoster vaccine recommended regardless of prior episode of zoster
Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)
No recommendation

Adapted from the Advisory Committee on Immunization Practices (ACIP) 2013 Adult Immunization Schedule. A summary of the adult immunization schedule vaccines and their primary indications, adverse events and contraindications can be found at: <a href="https://www.cdc.gov/vaccines/schedules/downloads/adult/mmwr-adult-schedule.pdf">www.cdc.gov/vaccines/schedules/downloads/adult/mmwr-adult-schedule.pdf</a>. For more detailed information on immunization of persons with HIV infection against influenza, pneumococcal disease, hepatitis B, human papillomavirus, varicella, and hepatitis A, see disease-specific sections in the text and in Table 1. For additional information on these and other vaccines (tetanus, diphtheria, pertussis, measles, mumps, rubella, and meningococcal disease), refer to recommendations of the ACIP at: <a href="https://www.cdc.gov/vaccines/pubs/acip-list.htm">www.cdc.gov/vaccines/pubs/acip-list.htm</a>.



<sup>\*</sup>IIV - Inactivated Influenza Vaccine. LAIV (live attenuated influenza vaccine) is not recommended for HIV-infected persons.